ELSEVIER

Carbohydrate Research 265 (1994) C9–C11

CARBOHYDRATE RESEARCH

CONTENTS

An electrochemical synthesis of methyl α-isomaltoside
C.H. Hamann, H. Polligkeit, P. Wolf (Oldenburg, Germany) and Z. Smiatacz (Gdańsk, Poland) 1

A short synthesis of β-xylobiosides
L. Ziser and S.G. Withers (Vancouver, British Columbia, Canada) 9

Differential chemical modification of substrate binding areas in porcine-pancreatic alpha-amylase by three regioisomeric photolabile ligands
J. Lehmann (Freiburg i. Br., Germany), G. Marchis-Mouren (Marseille, France), E. Schiltz and M. Schmidt-Schuchardt (Freiburg i. Br., Germany) 19

Di-D-fructose dianhydrides from the pyrolysis of inulin
A.E. Blize, M. Manley-Harris and G.N. Richards (Missoula, MT, USA) 31

Structural determination of the capsular antigen of *Escherichia coli* O8:K25:H9
A.N. Anderson, L.A.S. Parolis and H. Parolis (Grahamstown, South Africa) 41

A novel mono-sulfated pentaglycosylceramide with the isoglobo-series core structure in rat kidney
K. Tadano-Aritomi, M. Okuda, I. Ishizuka (Tokyo, Japan), H. Kubo (Kyoto, Japan) and P. Ireland (Manchester, United Kingdom) 49

Structural features of cell wall polysaccharides from the cotyledons of mung bean *Vigna radiata*
J. Gooneratne (Colombo, Sri Lanka), P.W. Needs, P. Ryden and R.R. Selvendran (Norwich, United Kingdom) 61

NMR assignment and conformational analysis of the antigenic capsular polysaccharide from *Streptococcus pneumoniae* type 9N in aqueous solution
T.J. Rutherford, C. Jones (Hertfordshire, United Kingdom), D.B. Davies (London, United Kingdom) and A.C. Elliott (Berkshire, United Kingdom). 79

Molecular recognition of antigenic polysaccharides: a conformational comparison of capsules from *Streptococcus pneumoniæ* serogroup 9
T.J. Rutherford, C. Jones (Hertfordshire, United Kingdom), D.B. Davies (London, United Kingdom) and A.C. Elliott (Berkshire, United Kingdom). 97

Structural studies of the enterotoxigenic *Escherichia coli* (ETEC) O153 O-antigenic polysaccharide
S. Ratnayake (Stockholm, Sweden), A. Weintraub (Huddinge, Sweden) and G. Widmalm (Stockholm, Sweden) 113

Structural studies of the *Shigella boydii* type 5 O-antigen polysaccharide
M.J. Albert (Dhaka, Bangladesh), T. Holme, B. Lindberg, J. Lindberg (Stockholm, Sweden), M. Mosihuzzaman, F. Qadri and M.M. Rahman (Dhaka, Bangladesh) 121

Synthesis of cyclohexakis- and cycloheptakis-(1 → 4)-(7-amino-6,7-dideoxy-α-D-*gluco*-heptopyranosyl), homoanalogues of 6-amino-6-deoxy-cyclomaltooligosaccharides
J. Defaye and A. Gadelle (Grenoble, France) 129

A convenient preparation of 5-benzyl ethers of D-gluco- and D-manno-furanose derivatives
H. Stępowska and A. Zamojski (Warsaw, Poland) 133

Synthesis of acylphosphonyl analogues of phosphatidyl-*myo*-inositol as inhibitors of phosphatidylinositol-specific phospholipase C
M.A. Alisi, M. Brufani, L. Filocamo, L. Cellai, M.A. Iannelli (Roma, Italy), M.C. Cesta and S. Lappa (Milano, Italy) 139

On the selectivity of stannylene-mediated alkylation and esterification of methyl 4,6-*O*-benzylidene α-D-glucopyranoside
D.J. Jenkins and B.V.L. Potter (Bath, United Kingdom) 145

A conformational study of the Smith degradation product of the *Klebsiella* K40 capsular polysaccharide by 1D NOESY and molecular mechanics calculations
R. Toffanin, M. Matulová, J. Bella (Trieste, Italy), D. Lamba (Rome, Italy), P. Cescutti, S. Paoletti and B.J. Kvam (Trieste, Italy) 151

Polymer-supported enzymic synthesis on a preparative scale
S. Köpper (Oldenburg, Germany) 161

NMR analysis of succinoglycans from different microbial sources: partial assignment of their ^{1}H and ^{13}C NMR spectra and location of the succinate and the acetate groups
M. Matulová, R. Toffanin, L. Navarini, R. Gilli, S. Paoletti and A. Cesàro (Trieste, Italy) . . . 167

Induced circular dichroism study of the aqueous solution complexation of cello-oligosaccharides and related polysaccharides with aromatic dyes
A.R. Engle, N. Purdie (Stillwater, OK, USA) and J.A. Hyatt (Kingsport, TN, USA) 181

Conformations and structure studies of sugar lactones. Part III. The composition and conformation of D-mannurono-γ-lactone in solution, and the structural analysis of its β anomer in the solid state
M.A. Shalaby, F.R. Fronczek, D. Vargas and E.S. Younathan (Baton Rouge, LA, USA). . . . 197

The configuration and conformation of di-D-fructose anhydride I. The crystal and molecular structure of 3,4,3′,4′-tetra-*O*-acetyl-6,6′-di(triphenylmethyl)-di-D-fructose anhydride I
M.A. Shalaby, F.R. Fronczek and E.S. Younathan (Baton Rouge, LA, USA) 207

The effect of stereochemistry upon carbohydrate hydration. A molecular dynamics simulation of β-D-galactopyranose and (α,β)-D-talopyranose
S.A. Galema (Groningen, Netherlands), E. Howard (La Plata, Argentina), J.B.F.N. Engberts (Groningen, Netherlands) and J.R. Grigera (La Plata, Argentina) 215

Facile synthesis of the sugar cores from phenylpropanoid glycosides
Z.-J. Li, H.-Q. Huang and M.-S. Cai (Beijing, China) 227

Preparation and crystal and molecular structure of 6-*O*-[(2*S*)-2,3-epoxypropyl]-1,2:3,4-di-*O*-isopropylidene-α-D-galactopyranose. Pyranoid ring conformation in 1,2:3,4-di-*O*-isopropylidene-galactopyranose and related systems
P. Köll, W. Saak, S. Pohl (Oldenburg, Germany), B. Steiner and M. Kožíš (Bratislava, Slovak Republic) 237

Difructose dianhydrides from sucrose and fructo-oligosaccharides and their use as building blocks for the preparation of amphiphiles, liquid crystals, and polymers
J.M. García Fernández, A. Gadelle and J. Defaye (Grenoble, France) 249

Regioselective chlorination of chitin with *N*-chlorosuccinimide-triphenylphosphine under homogeneous conditions in lithium chloride–*N*,*N*-dimethylacetamide
M. Sakamoto, H. Tseng and K.-i. Furuhata (Tokyo, Japan) 271

Effects of added galacto- and gluco-mannans on the NMR spectra of ^{133}Cs ions in kappa-carrageenan gels
L. Piculell, W. Zhang (Lund, Sweden), T. Turquois (Nantes, France), C. Rochas (Saint Martin d'Hères, France), F.-R. Taravel (Grenoble, France) and P.A. Williams (Deeside, United Kingdom) 281

Notes

The crystal and molecular structure of 4-cyanophenyl and 4-nitrophenyl β-D-xylopyranosides
J.Y. Le Questel, N. Mouhous-Riou and S. Pérez (Nantes, France) 291

Synthesis of 5-fluorouridine 5′-diphosphate galactose from 5-fluorouridine by chemical phosphorylation and microbial uridylyl transfer
K.-i. Fujita, A. Matsukawa, K. Shibata, T. Tanaka, M. Taniguchi and S. Oi (Osaka, Japan) . . 299

NMR spectroscopic investigation of oligoglucuronates prepared by enzymic hydrolysis of a (1 → 4)-β-D-glucuronan
L. Dantas (Grenoble, France), J. Courtois, B. Courtois, J.-P. Seguin (Amiens, France), C. Gey and A. Heyraud (Grenoble, France) 303

Preparation of 2,3,6,2′,3′,4′,6′-hepta-*O*-acetyl-maltose/cellobiose by enzymatic hydrolysis of maltose/cellobiose octaacetate
G.-T. Ong, K.-Y. Chang, S.-H. Wu and K.-T. Wang (Taipei, Taiwan) 311

Structure of the exopolysaccharide of *Pseudomonas stutzeri* strain ATCC 17588
S.F. Osman, W.F. Fett and R.L. Dudley (Philadelphia, PA, USA) 319

Preparation of *N*-acetylchitooligosaccharides by hydrolysis of chitosan with chitinase followed by *N*-acetylation
S.-i. Aiba (Tsukuba, Japan) 323

Announcements C1